

UC Berkeley

UC Berkeley Previously Published Works

Title

Photoinitiated oxidative addition of CF₃I to gold(I) and facile aryl-CF₃ reductive elimination.

Permalink

<https://escholarship.org/uc/item/9rr6x0c5>

Journal

Journal of the American Chemical Society, 136(21)

ISSN

0002-7863

Authors

Winston, Matthew S
Wolf, William J
Toste, F Dean

Publication Date

2014-05-01

DOI

10.1021/ja503974x

Peer reviewed


Photoinitiated Oxidative Addition of CF₃I to Gold(I) and Facile Aryl-CF₃ Reductive Elimination

Matthew S. Winston, William J. Wolf, and F. Dean Toste*

Department of Chemistry, University of California, Berkeley, California 94720, United States

Supporting Information

ABSTRACT: Herein we report the mechanism of oxidative addition of CF₃I to Au(I), and remarkably fast C_{aryl}-CF₃ bond reductive elimination from Au(III) cations. CF₃I undergoes a fast, formal oxidative addition to R₃PAuR' (R = Cy, R' = 3,5-F₂-C₆H₄, 4-F-C₆H₄, C₆H₅, 4-Me-C₆H₄, 4-MeO-C₆H₄, Me; R = Ph, R' = 4-F-C₆H₄, 4-Me-C₆H₄). When R' = aryl, complexes of the type R₃PAu(aryl)(CF₃)I can be isolated and characterized. Mechanistic studies suggest that near-ultraviolet light (λ_{max} = 313 nm) photoinitiates a radical chain reaction by exciting CF₃I. Complexes supported by PPh₃ undergo reversible phosphine dissociation at 110 °C to generate a three-coordinate intermediate that undergoes slow reductive elimination. These processes are quantitative and heavily favor C_{aryl}-I reductive elimination over C_{aryl}-CF₃ reductive elimination. Silver-mediated halide abstraction from all complexes of the type R₃PAu(aryl)(CF₃)I results in quantitative formation of Ar-CF₃ in less than 1 min at temperatures as low as -10 °C.




INTRODUCTION

Reports of organogold complexes undergoing redox processes are typically limited to slow oxidative additions and reductive eliminations.^{1,2} However, organogold complexes are not necessarily unreactive; we recently showed that diaryl Au(III) complexes undergo remarkably fast aryl-aryl reductive elimination at temperatures as low as -50 °C.³ These recent findings from our group, as well those established by Vicente,⁴ Hashmi,⁵ and Lloyd-Jones,⁶ suggest that the barrier for challenging reductive eliminations might be substantially diminished at Au(III). C_{aryl}-CF₃ bond reductive elimination is typically a slow process requiring elevated temperatures and long reaction times, due to ground state stabilization afforded by exceptionally strong bonding between transition metals and CF₃ ligands.⁷ For instance, (dppbz)Pd(2-Me-C₆H₄)(CF₃) (dppbz = 1,2-bis(diphenylphosphino)benzene) is stable at 130 °C for 3 days,⁸ while (dppp)Pd(Ph)(CF₃) (dppp = 1,3-diphenylphosphinopropane) and (dppe)Pd(Ph)(CF₃) (dppe = 1,2-diphenylphosphinoethane) yield only 10% PhCF₃ after 3 days at 145 °C.⁹ Reductive eliminations at temperatures between 50 and 80 °C can be achieved at Pd(II) by employing bulky ligands, such as Xantphos¹⁰ and Brettphos.¹¹ Notably, while aryl-CF₃ reductive eliminations from Pd(IV) often require similarly high temperatures,^{12a} Sanford has shown that they can occur at temperatures as low as 23 °C over 1 h.^{12b} Despite advances in catalytic trifluoromethylation, C_{aryl}-CF₃ reductive elimination still remains a challenging step. Given the importance of trifluoromethylated arenes in pharmaceuticals and agrochemicals,¹³ we were prompted to investigate potentially low-barrier C_{aryl}-CF₃ bond reductive elimination at Au(III).

To access complexes of the type R₃PAu(aryl)(CF₃)I, we were drawn to Puddephatt's report of the oxidative addition of CF₃I to Me₃PAuMe to afford *cis/trans* mixtures of Me₃PAuMe₂(CF₃) and Me₃PAuI.¹⁴ In one case, Me₃PAu(Me)-(CF₃)I was obtained exclusively, but its preparation could not be reproduced by the authors. Because reaction times varied from 5 min to 1 day, and rates dramatically slowed in the presence of galvinoxyl, the authors concluded that a free-radical chain mechanism was operative, with •CF₃ as the propagating species.

RESULTS AND DISCUSSION

Prior investigations by our group revealed that oxidation of Ph₃PAu(4-F-C₆H₄) rapidly generates 4,4'-difluorobiphenyl through a mechanism involving aryl group transfer.³ However, the use of the bulkier PCy₃ prevents transfer of the arene ligand, instead resulting in clean, rapid oxidation of Cy₃PAu(4-F-C₆H₄) (**1a**) to the isolable Au(III) complex *cis*-(Cy₃P)Au(4-F-C₆H₄)Cl₂ (**2**) (eq 1).¹⁵ Therefore, we began our inves-



tigations of Au(I) oxidation by CF₃I using **1a**, with the fluorinated arene ligand also providing a convenient ¹⁹F NMR handle. Treatment of **1a** in CD₂Cl₂ with CF₃I (25 equiv) afforded the product of formal CF₃I oxidative addition **3a** in 1 h in good yield (eq 2 and Table 1). Both the CF₃ and PCy₃

Received: April 21, 2014

Published: May 16, 2014




Table 1. Photoinitiated Oxidative Addition of CF₃I to Electronically Diverse Au(I) Aryl Complexes 1a–1f, 11a, and 11b

$\text{R}_3\text{PAuAr} \xrightarrow[\text{CD}_2\text{Cl}_2]{\text{CF}_3\text{I (25 equiv), } h\nu (313 \text{ nm})} \text{R}_3\text{P} \rightarrow \text{Au} \begin{array}{l} \text{CF}_3 \\ \text{I} \\ \text{Ar} \end{array}$				
Au(I) reactant	PR ₃	Ar	product	yield (%)
1a	PCy ₃	4-F-C ₆ H ₄	3a	64
1b	PCy ₃	3,5-F ₂ -C ₆ H ₃	3b	44
1c	PCy ₃	C ₆ H ₅	3c	59
1d	PCy ₃	4-Me-C ₆ H ₄	3d	38
1e	PCy ₃	4-MeO-C ₆ H ₄	3e	44
1f	PCy ₃	2-Me-C ₆ H ₄	—	NR
11a	PPh ₃	4-F-C ₆ H ₄	12a	71
11b	PPh ₃	4-Me-C ₆ H ₄	12b	63

ligands (doublet at $\delta = -24.5$ and quartet at $\delta = 25.6$ in the ¹⁹F and ³¹P NMR spectra, respectively) provide diagnostic NMR signals (Table 2). The substantial coupling (³J_{P–F} = 63 Hz) between fluorine and phosphorus are characteristic of a *trans* relationship between the CF₃ and phosphine ligands.¹⁴ X-ray analysis of crystals of **3a** confirmed this stereochemical relationship around the square planar Au(III) (Figure 1A); other than the homoleptic anion [Au(CF₃)₄][–],¹⁶ complex **3a** contains a rare example of a crystallographically characterized Au(III)–CF₃ bond. Complex **3a** is not only stable to air and water but can be purified by column chromatography as well.

Mechanism of Oxidative Addition of CF₃I. The reaction of **1a** and CF₃I represents a rare oxidation of Au(I) to Au(III) that directly installs potentially reactive Au(III)–carbon bonds.¹ During our attempts to monitor the oxidative addition by ¹⁹F NMR, we found that no reaction occurred when the reaction mixture was placed inside the dark NMR spectrometer. However, when the reaction mixture was exposed to ambient fluorescent light for 5 min, the formation of **3a** was detected (~20%). Given the reliance of numerous methods on CF₃I as a trifluoromethyl source,¹⁷ we investigated its photochemical reactivity. Actinometry experiments were carried out to determine the overall quantum yield, using the Norrish II

fragmentation of valerophenone as a standard.¹⁸ The oxidative addition of CF₃I to **1a** was complete after 20 s of irradiation by a Hg vapor lamp (2 mM aq. K₂CrO₄ optical filter; transmittance $\lambda_{\text{max}} = 313 \text{ nm}$), while the fragmentation of valerophenone ($\Phi = 1$) took place over 24 h under identical conditions. This rate difference, in addition to the ability of ambient light to bring the reaction to full conversion over variable reaction times (between 15 min and 1 h), supports a radical chain reaction as the mechanism of Au(I) oxidation by CF₃I.

The reaction of excess CF₃I and **1a** is also fast in THF, but the conversion is never greater than 65% (52% yield of **3a**), even when irradiated by a Hg vapor lamp for 1 h (*vide infra*). Notably, an excess of fluorocarbon (HCF₃) is generated in THF, regardless of the light source (only DCF₃ is formed when THF-d₈ is used). GC-MS analysis of reaction mixtures reveals several products of THF oxidation, likely formed by H• abstraction by •CF₃.

Several control experiments, using HCF₃ production relative to a standard as a probe to detect •CF₃ generation, support the involvement of Au(I) during the initiation of the chain reaction. The UV absorption of CF₃I is centered at 270 nm but tails beyond 350 nm.¹⁹ When irradiated at 313 nm, CF₃I undergoes fast, reversible C–I bond homolysis. However, in the absence of **1a**, only negligible amounts of HCF₃ are observed when THF solutions of CF₃I are irradiated for 30 min, indicating that carbon/iodine radical recombination is substantially faster than H• abstraction from THF. Similarly insignificant quantities of HCF₃ are observed when 20 equiv (relative to CF₃I) of the H• donors 1,4-cyclohexadiene, 9,10-dihydroanthracene, or triphenylmethane are added (Figure 2A). Additionally, Cy₃PAu(2-(CH₂CH=CH₂)C₆H₄) (**4**), containing a pendent olefin to either capture a putative Au(II) intermediate and/or •CF₃, is fully consumed upon irradiation in the presence of excess CF₃I (Figure 2B). This oxidation affords multiple Au(III) products of indiscriminate •CF₃ addition to the terminal olefin and gold atom (and HCF₃ when THF is used as solvent) (see SI). Because 2-allylbromobenzene (**5**) does not react with CF₃I when irradiated under similar conditions (no HCF₃ is observed after 5 min, and less than 2% after 30 min), we conclude that the Au(I) aryl complex is necessary for chain initiation. These results are also consistent with an initiation mechanism involving [CF₃I]^{•–}, which generates iodide and •CF₃ following C–I bond homolysis.

We envisioned two possible initiation mechanisms for generating •CF₃ as a propagating species from [CF₃I]^{•–} in a chain reaction: (1) initial photoexcitation of **6** followed by electron transfer to CF₃I, or (2) initial photoexcitation of CF₃I followed by electron transfer from **6** (Scheme 1).

Table 2. ³¹P{¹H} and ¹⁹F NMR Data for Complexes 3a–3e, 12a, and 12b

$\text{R}_3\text{P} \rightarrow \text{Au} \begin{array}{l} \text{CF}_3 \\ \text{I} \\ \text{Ar} \end{array}$				
complex	PR ₃	Ar	δ ³¹ P{ ¹ H} (ppm)	δ ¹⁹ F (ppm)
3a	PCy ₃	4-F-C ₆ H ₄	25.6 (q, ³ J _{P–F} = 63 Hz)	–24.5 (d, ³ J _{P–F} = 63 Hz)
3b	PCy ₃	3,5-F ₂ -C ₆ H ₃	26.1 (q, ³ J _{P–F} = 63 Hz)	–22.0 (d, ³ J _{P–F} = 64 Hz)
3c	PCy ₃	C ₆ H ₅	25.5 (q, ³ J _{P–F} = 62 Hz)	–22.7 (d, ³ J _{P–F} = 62 Hz)
3d	PCy ₃	4-Me-C ₆ H ₄	25.5 (q, ³ J _{P–F} = 62 Hz)	–23.6 (d, ³ J _{P–F} = 62 Hz)
3e	PCy ₃	4-MeO-C ₆ H ₄	23.3 (q, ³ J _{P–F} = 63 Hz)	–20.6 (d, ³ J _{P–F} = 64 Hz)
12a	PPh ₃	4-F-C ₆ H ₄	20.0 (q, ³ J _{P–F} = 68 Hz)	–21.0 (d, ³ J _{P–F} = 68 Hz)
12b	PPh ₃	4-Me-C ₆ H ₄	20.4 (q, ³ J _{P–F} = 67 Hz)	–21.3 (d, ³ J _{P–F} = 67 Hz)




Figure 1. (A–F) Thermal ellipsoid representations of 3a–3d, 12a, and 12b at the 50% probability level. Hydrogens have been omitted for clarity. Atoms are color-coded: gray (carbon), yellow (fluorine), gold (gold), purple (iodine), orange (phosphorus). See Supporting Information (SI) for bond lengths and angles.





Figure 2. Control experiments to assess involvement of Au(I) in the initiation of the radical chain mechanism. (A) Irradiation of CF_3I solutions containing H^\bullet donors to detect CF_3H in the absence of gold. (B) Radical trapping using an olefin with and without a pendant gold center.

Au(I) aryl complexes are well-known chromophores, and their photophysical properties have been investigated previously.²⁰ While **1a** absorbs weakly above 310 nm (the cutoff for many laboratory fluorescent lamps¹⁹), excitation at 320 nm ($\epsilon = 37 \text{ M}^{-1} \text{ cm}^{-1}$) results in a weak, broad luminescence from 340 to 460 nm, classified as fluorescence based on the lifetime of excited species **1a*** (<10 ns, quantum yield of fluorescence = 0.03).²¹ Despite the short lifetime of **1a***, CF_3I effectively quenches its fluorescence (Stern–Volmer quenching constant $K_{\text{SV}} = 30 \text{ M}^{-1}$, Figure 3). Although this energy transfer could conceivably generate $\cdot\text{CF}_3$ and initiate a chain reaction (mechanism 1, Scheme 1), when CF_3I is removed from

Scheme 1. Possible Initiation Mechanisms Involving Photoexcitation of Either Au(I) Complex **6** (Mechanism 1) or CF_3I (Mechanism 2)



fluorimetry samples under vacuum, fluorescence is restored to the same intensity prior to introduction of the gas, indicating that consumption of Au(I) has not occurred.

Surprisingly, fluorescence quenching by the Au(III) complex **3a** is more than 2 orders of magnitude more effective ($K_{\text{SV}} = 4270 \text{ M}^{-1}$) than quenching by CF_3I (Figure 3). If propagating species terminate frequently, some critical concentration of Au(III) product exists that may impede productive energy transfer from an excited species, halting reinitiation of the chain reaction.

In light of Puddephatt's report, Au(I) alkyl complexes, such as Me_3PAuMe , clearly react with CF_3I .¹⁴ However, there is no mention of the dependence of light on this process, although if the reaction is photoinitiated, mechanism 1 would seem especially unlikely given the absence of a chromophoric aryl ligand in Puddephatt's examples. To test this hypothesis, we irradiated Cy_3PAuMe (**9**) in the presence of CF_3I (Scheme 2).





Figure 3. Stern–Volmer plots of fluorescence quenching of **1a** by different concentrations of CF_3I (blue boxes) and Au(III) complex **3a** (blue triangles) in CH_2Cl_2 . Concentrations of Au(III) are in mol/L and CF_3I concentrations are in mmol/L.


Scheme 2. Photochemical Oxidative Addition of CF_3I to **9 in CD_2Cl_2 and Spontaneous Reductive Elimination of CH_3I**



While **9** does not absorb above 300 nm (see SI), the reaction is quantitative in CD_2Cl_2 when irradiated with ambient light, and *does not proceed in the dark*. The oxidized product is unobservable, eliminating CH_3I to generate $\text{Cy}_3\text{PAuCF}_3$ at room temperature.^{22,23} In THF, the reaction generates excess HCF in THF_3 , presumably also from solvent H^\bullet abstraction by $^\bullet\text{CF}_3$.


If initiation mechanism 2 is operative, then $^\bullet\text{CF}_3$ could be generated by irradiating CF_3I solutions containing electron donors other than Au(I) , such as phosphines (Scheme 3).²⁴

Scheme 3. Photochemical Oxidation of Trialkylphosphines by CF_3I



Indeed, irradiation of PMe_3 or PCy_3 in the presence of CF_3I results in formation of $[\text{Me}_3\text{P-CF}_3]\text{I}$ (**10a**, $^2J_{\text{P-F}} = 63 \text{ Hz}$) or $[\text{Cy}_3\text{P-CF}_3]\text{I}$ (**10b**, $^2J_{\text{P-F}} = 42 \text{ Hz}$);²⁵ neither reaction proceeds in the dark. Consistent with quenching of $[\text{CF}_3\text{I}]^*$ by Au(III) , the oxidation of PCy_3 in THF stalls at roughly 45% conversion (by ^{31}P NMR) in the presence of 25 mol % Au(III) complex **3a**.

PPh_3 does not react with CF_3I (eq 3), presumably due to its lower oxidation potential relative to PMe_3 and PCy_3 . When PCy_3 and PPh_3 are irradiated *together* with CF_3I , only PCy_3 is consumed, suggesting that PPh_3 neither initiates the chain nor reacts with $^\bullet\text{CF}_3$ during propagation. Contrary to our initial hypothesis that bulky phosphine ligands prevent aryl group transfer upon Au(I) oxidation, we found that $\text{Ph}_3\text{PAu(4-F-C}_6\text{H}_4)$ (**11a**) undergoes quantitative photoinitiated reaction with CF_3I in CD_2Cl_2 to generate **12a** (eq 4). Since PPh_3 is




unreactive toward CF_3I , oxidation of **11a** cannot be initiated by small amounts of dissociated PPh_3 (we cannot disprove the analogous mechanism for PCy_3 -supported complex **1a**.) Complex **12a** was characterized by X-ray crystallography and shown to be isostructural to **3a** (Figure 1B).

On the basis of these results, we propose that while photoexcited $[\text{CF}_3\text{I}]^*$ undergoes rapid C–I bond homolysis and recombination, it also oxidizes Au(I) aryl and alkyl complexes by accepting electrons into a low-lying SOMO to generate radical anion $[\text{CF}_3\text{I}]^{\bullet-}$ (mechanism 2, Scheme 1). Homolysis of the C–I bond of $[\text{CF}_3\text{I}]^{\bullet-}$ generates iodide and $^\bullet\text{CF}_3$, which oxidizes $(\text{R}_3\text{P})\text{AuR}'$ (**6**) to Au(II) intermediate **7**. Iodine atom abstraction of CF_3I by **7** affords Au(III) complex **8** and regenerates $^\bullet\text{CF}_3$. In THF, oxidation of **6** by $^\bullet\text{CF}_3$ is competitive with solvent H^\bullet abstraction to make HCF_3 and terminate the radical chain. At sufficiently high concentrations, the Au(III) product (**8**) quenches $[\text{CF}_3\text{I}]^*$ before it can reinitiate the radical chain reaction.

Promisingly, the photoinitiated oxidative addition of CF_3I is general for electronically diverse complexes of the type $\text{Cy}_3\text{PAu(aryl)}$ (Tables 1 and 2). The resulting Au(III) products (see Figure 1 for their crystallographic analyses) can be purified by chromatography on silica. Complex **1b** (aryl = 3,5- $\text{F}_2\text{-C}_6\text{H}_3$), which is more electron-deficient than **1a** (aryl = 4- $\text{F-C}_6\text{H}_4$), reacts smoothly with CF_3I to afford **3b**. While complexes with more electron-rich ligands such as **1c** (aryl = C_6H_5) and **1d** (aryl = 4- $\text{Me-C}_6\text{H}_4$) also react with CF_3I to afford **3c** and **3d**, respectively, the most electron-rich complex **1e** (aryl = 4- $\text{MeO-C}_6\text{H}_4$) decomposes to Au nano particles and several CF_3 -containing Au(III) complexes in solution and solid state (no products of $\text{C}_{\text{aryl}}\text{-I}$ or $\text{C}_{\text{aryl}}\text{-CF}_3$ reductive elimination can be detected). Au(III) product **3e** is detectable, however, and its decomposition can be slowed substantially by addition of MeCN upon concentration of the reaction, allowing its solution-state characterization. The mechanism of decomposition has not yet been identified, although we speculate that the electron-rich arene may encourage PCy_3 dissociation at room temperature and subsequent aryl group transfer.

The complex **1f** (aryl = 2- $\text{Me-C}_6\text{H}_4$) does not react with CF_3I at all, suggesting that CF_3I oxidative addition is sensitive to the sterics of the aryl ligand and that relaxation of $[\text{CF}_3\text{I}]^*$ is faster than oxidation of the metal center to initiate the radical chain. Unsurprisingly, no HCF_3 is observed when **1f** is irradiated in THF for 20 min.

Reductive Elimination from Au(III) Complexes. We next probed $\text{C}_{\text{aryl}}\text{-CF}_3$ reductive eliminations from Au(III) . To our surprise, **12a** undergoes quantitative $\text{C}_{\text{aryl}}\text{-I}$ reductive elimination in toluene- d_8 at 110 °C to afford 4-fluoriodobenzene and $\text{Ph}_3\text{PAuCF}_3$ over 20 min (Scheme 4).²² No 4-fluoro(trifluoromethyl)benzene is observed by ^{19}F NMR or GC. This process is highly sensitive to free phosphine, stalling completely in the presence of PPh_3 (0.1 or 1.0 equiv) at 110 °C for 12 h. Treatment of **12a** with $\text{PPh}_3\text{-}d_{15}$ at room temperature

Scheme 4. Behavior of Au(III) Complex 12a and 12b in the Presence of Free PPh₃ and at Elevated Temperatures


results in immediate formation of **12a-d₁₅**, presumably via an associative process.^{2a-c}

More electron-rich aryl ligands, such as 4-methylphenyl (**12b**), do not significantly affect the relative rates of C_{aryl}-I and C_{aryl}-CF₃ reductive elimination (Scheme 4). At 110 °C, complex **12b** undergoes mostly C_{aryl}-I reductive elimination within 10 min to afford 4-methyliodobenzene.²⁶ Both C_{aryl}-I and C_{aryl}-CF₃ reductive eliminations are also completely inhibited in the presence of PPh₃ (0.1 or 1.0 equiv), while PPh₃-d₁₅ reacts immediately at room temperature to afford **12b-d₁₅**, also via associative ligand exchange. These observations are consistent with a mechanism involving highly reversible PPh₃ dissociation from **12a** and **12b**, followed by slow C_{aryl}-I reductive elimination from **13a** or **13b**, respectively.

Clearly, the behaviors of **12a** and **12b** are similar to Au(III)alkyl complexes studied by Kochi, which not only reductively eliminate C_{alkyl}-C_{alkyl} bonds between 70 and 100 °C via a dissociative mechanism but also undergo associative ligand exchange at ambient temperature with excess phosphine.^{2a-c} Unsurprisingly, analogous PCy₃-stabilized complexes **3a** and **3d** are stable at 110 °C for at least 12 h, presumably due to the greater σ-donating ability of PCy₃ relative to PPh₃. Phosphine exchange with excess P(*n*-Bu)₃, PBn₃, or PCy₃ does not occur even at these temperatures, precluding not only the lower-barrier associative exchange mechanism observed with the PPh₃-supported systems (attributed to the larger cone angle of PCy₃ relative to PPh₃), but also PCy₃ dissociation to form a three-coordinate complex.

Because C_{aryl}-I reductive elimination is significantly faster than C_{aryl}-CF₃ reductive elimination, a cycle for gold-catalyzed trifluoromethylation must necessarily involve iodide abstraction from the Au(III) product of CF₃I oxidative addition. Despite the apparent kinetic stabilities of the Au(III) complexes **3a–3e**, **12a**, and **12b**, they all undergo quantitative C_{aryl}-CF₃ reductive elimination in less than 1 min upon treatment with AgSbF₆ at room temperature.


To consider the effects of the phosphine ligand on the silver-mediated C_{aryl}-CF₃ reductive elimination of Au(III), we used variable-temperature NMR to follow the reductive elimination from **3a** and **12a** in the presence of AgSbF₆. PCy₃-substituted complex **3a** undergoes very fast (quantitative conversion in less than 1 min) C_{aryl}-CF₃ reductive elimination at −10 °C, while the analogous PPh₃-stabilized **12a** reacts similarly fast at room temperature (eq 5). At lower temperatures, several bridging species (most likely dimers) are observed by ¹⁹F NMR upon halide abstraction in both cases. If C_{aryl}-CF₃ bond reductive elimination can only occur from a monomeric three-coordinate intermediate, then **12a** might be expected to undergo slower



reductive elimination due to slower dimer dissociation and/or a dimer–monomer equilibrium that more favors the dimer, based on the smaller cone angle and weaker σ-donation of PPh₃ relative to PCy₃.

CONCLUSION

These results reported herein support the oxidative addition of CF₃I to Au(I) via a photoinitiated chain reaction. The reactions are fast at room temperature for both Au(I) aryl and alkyl complexes. Aryl-CF₃ reductive elimination is typically a high-barrier process but occurs in seconds at room temperature from a Au(III) cation. The Au(I)aryl species may be regenerated via one of the numerous transmetalation strategies available involving carbon nucleophiles.²⁷ For instance, excess (4-F-C₆H₄)SnMe₃ (10 equiv) undergoes fast, quantitative transmetalation with [Cy₃PAu]SbF₆ at room temperature to afford **1a**, thereby closing a hypothetical catalytic cycle based on the three elementary steps shown in Scheme 5. Silver-free halide

Scheme 5. Oxidation of **1a**, Aryl-CF₃ Reductive Elimination, and Regeneration of **1a** Supports the Feasibility of a Mild, Catalytic Trifluoromethylation

abstraction from Au(III) complexes could conceivably enable a practical and mild cycle for gold-catalyzed trifluoromethylation of aryl nucleophiles, although deleterious reactions between starting material and metalloradical intermediates and •CF₃ must be mitigated, as well as competitive aryl–aryl homocoupling.

While we initially set out to probe C_{aryl}-CF₃ reductive elimination at Au(III), we also explored the oxidative addition of CF₃I to Au(I), a process with potential implications beyond gold chemistry. The possibility of photoinitiated oxidation of transition metals or main group elements by CF₃I should not be discounted in methods employing this reagent as a trifluoromethyl source, particularly since ambient fluorescent laboratory lighting is sufficient to initiate a chain in the presence of a suitable reductant. The results presented also suggest that substrate photoexcitation may provide a low-barrier avenue to kinetically challenging oxidative additions by Au(I), providing access to potentially reactive Au(III) complexes.²⁸

ASSOCIATED CONTENT

Supporting Information

Experimental details, characterization data, and crystallographic information (cif). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

E-mail: fdtoste@berkeley.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We gratefully acknowledge Professor Robert G. Bergman for helpful discussions, David Tatum and Professor Kenneth N. Raymond for use of fluorimetry equipment, Professor Felix Fischer for access to a Hanovia lamp, and Mercedes Taylor and the Chemistry 208 course at UC-Berkeley for assistance with the X-ray structure of **3b**. This work was generously funded by the NIHGMs (RO1 GM073932), an NIH fellowship to M. S. W. (F32 GM103238-02), and an NSF fellowship to W.J.W. (DGE 1106400).

■ REFERENCES

- (1) For oxidative addition: (a) Johnson, A.; Puddephatt, R. J. *Inorg. Nucl. Chem. Lett.* **1973**, *9*, 1175. (b) Tamaki, A.; Kochi, J. K. *J. Organomet. Chem.* **1973**, *64*, 411. (c) Johnson, A.; Puddephatt, R. J. *J. Organomet. Chem.* **1975**, *85*, 115. (d) Shiotani, A.; Schmidbaur, H. *J. Organomet. Chem.* **1972**, *37*, C24. (e) Tamaki, A.; Magennis, S. A.; Kochi, J. K. *J. Am. Chem. Soc.* **1973**, *95*, 6487. (f) Fackler, J. P., Jr. *Polyhedron* **1997**, *16*, 1.
- (2) For reductive elimination: (a) Tamaki, A.; Kochi, J. K. *J. Organomet. Chem.* **1974**, *64*, 411. (b) Tamaki, A.; Magennis, S. A.; Kochi, J. K. *J. Am. Chem. Soc.* **1974**, *96*, 6140. (b) Komiya, S.; Albright, T. A.; Hoffmann, R.; Kochi, J. K. *J. Am. Chem. Soc.* **1976**, *98*, 7255. (c) Komiya, S.; Kochi, J. K. *J. Am. Chem. Soc.* **1976**, *98*, 7599. (d) Kuch, P. L.; Tobias, R. S. *J. Organomet. Chem.* **1976**, *122*, 429.
- (3) Wolf, W. J.; Winston, M. S.; Toste, F. D. *Nat. Chem.* **2014**, *6*, 159.
- (4) (a) Vicente, J.; Bermudez, M. D.; Escibano, J. *Organometallics* **1991**, *10*, 3380. (b) Vicente, J.; Bermudez, M. D.; Carrion, F. *Inorg. Chim. Acta* **1994**, *220*, 1. (c) Vicente, J.; Bermudez, M. D.; Carrion, F. J.; Jones, P. G. *Chem. Ber.* **1996**, *196*, 1395.
- (5) Hashmi, A. S. K.; Blanco, M. C.; Fischer, D.; Bats, J. W. *Eur. J. Org. Chem.* **2006**, 1387.
- (6) (a) Ball, L. T.; Lloyd-Jones, G. C.; Russell, C. A. *Science* **2012**, *337*, 1644. (b) Ball, L. T.; Lloyd-Jones, G. C.; Russell, C. A. *J. Am. Chem. Soc.* **2014**, *136*, 254.
- (7) (a) Hughes, R. P. *Adv. Organomet. Chem.* **1990**, *31*, 183. (b) Morrison, J. A. *Adv. Organomet. Chem.* **1993**, *35*, 311.
- (8) Cullin, D. A.; Hartwig, J. F. *Organometallics* **2004**, *23*, 3398.
- (9) Grushin, V. V.; Marshall, W. J. *J. Am. Chem. Soc.* **2006**, *128*, 4632.
- (10) Grushin, V. V.; Marshall, W. J. *J. Am. Chem. Soc.* **2006**, *128*, 12644.
- (11) Cho, E. J.; Senecal, T. D.; Kinzel, T.; Zhang, Y.; Watson, D. A.; Buchwald, S. L. *Science* **2010**, *328*, 1679.
- (12) (a) Ball, N. D.; Kampf, J. W.; Sanford, M. S. *J. Am. Chem. Soc.* **2010**, *132*, 2878. (b) Ball, N. D.; Gary, J. B.; Ye, Y.; Sanford, M. S. *J. Am. Chem. Soc.* **2011**, *133*, 7577.
- (13) Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.* **2011**, *111*, 4475.
- (14) Johnson, A.; Puddephatt, R. J. *Chem. Soc., Dalton Trans.* **1976**, 1360.
- (15) Nevado and co-workers have reported the synthesis of complexes of the type *cis*-Ph₃PAu(aryl)Cl₂, where aryl = C₆F₅ or 2,4,6-F₃-C₆H₂. Because of the electron-deficient aryl ligands, transmetalation between Au(I) and Au(III) is slow enough to allow full oxidation by PhICl₂. See: (a) Hofer, M.; Nevado, C. *Tetrahedron* **2013**, *69*, 5751. (b) Hofer, M.; Gomez-Bengoa, E.; Nevado, C. *Organometallics* **2014**, *33*, 1328.
- (16) Martínez-Salvador, S.; Falvello, L. R.; Martín, A.; Manjón, B. *Chem.—Eur. J.* **2013**, *19*, 14540.
- (17) (a) Nagib, D. A.; Scott, M. E.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2009**, *131*, 10875. (b) Kino, T.; Nagase, Y.; Ohtsuka, Y.; Yamamoto, K.; Uraguchi, D.; Tokuhisa, K.; Yamakawa, T. *J. Fluorine Chem.* **2010**, *131*, 98. (c) Ye, Y.; Sanford, M. S. *J. Am. Chem. Soc.* **2012**, *134*, 9034.
- (18) Wagner, P. J.; Kemppainen, A. E. *J. Am. Chem. Soc.* **1972**, *94*, 7495.
- (19) Nyden, M. R. Photodegradation of CF₃I. In *Fire Suppression System Performance of Alternative Agents in Aircraft Engine and Dry Bay Laboratory Simulations*; Gann, R. G., Ed.; National Institute of Standards and Technology: Washington, D.C., 1995; pp 77–95.
- (20) For some recent examples, see: (a) Partyka, D. V.; Zeller, M.; Hunter, A. D.; Gray, T. G. *Angew. Chem., Int. Ed.* **2006**, *45*, 8188. (b) Partyka, D. V.; Esswein, A. J.; Zeller, M.; Hunter, A. D.; Gray, T. G. *Organometallics* **2007**, *26*, 3279. (c) Vogt, R. A.; Gray, T. G.; Crespo-Hernández, C. E. *J. Am. Chem. Soc.* **2012**, *134*, 14808. (d) Visbal, R.; Ospino, I.; López-de-Luzuriaga, J. M.; Laguna, A.; Gimeno, M. C. *J. Am. Chem. Soc.* **2013**, *135*, 4712. (e) Crespo, O.; Díez-Gil, C.; Jones, P. G.; Laguna, A.; Ospino, I.; Tapias, J.; Villacampa, M. D.; Visbal, R. *Dalton Trans.* **2013**, *42*, 8298. (f) Monzittu, F. M.; Fernández-Moreira, V.; Lippolis, V.; Arca, M.; Laguna, A.; Gimeno, M. C. *Dalton Trans.* **2014**, *43*, 6212.
- (21) The excited state lifetime of **1a*** is shorter than the lower limit of detection of our fluorimeter (<10 ns).
- (22) Cy₃PAuCF₃ and Ph₃PAuCF₃ were previously synthesized by the reaction of R₃PAuOCH(CF₃)₂ with TMS-CF₃. See: Usui, T.; Noma, J.; Hirano, M.; Komiya, S. *Inorg. Chim. Acta* **2000**, *309*, 151.
- (23) Due to the thermal stability of complexes **3a**, **3d**, **12a**, and **12b**, we favor an outer-sphere reductive elimination of CH₃I. Johnson and Puddephatt proposed a similar mechanism; see reference 1c.
- (24) Buckler, S. A.; Doll, L.; Lind, F. K.; Epstein, M. J. *Org. Chem.* **1962**, *27*, 794.
- (25) The unusually small ²J_{P-F} for [Cy₃P-CF₃]**1** (**10b**) may be a result of geometric distortions at phosphorus, which nearly planarize the bulky cyclohexyl ligands and reduce s contributions to the P-CF₃ bond.
- (26) Upon full conversion of **12b**, we detect roughly 2% of 4-methyl-1-trifluoromethylbenzene, the product of C_{aryl}-CF₃ reductive elimination from **13b**, by ¹⁹F NMR and GC.
- (27) See procedures in refs 1–4 and 19, as well as: (a) Hashmi, A. S. K.; Ramamurthi, T. D.; Rominger, F. J. *Organomet. Chem.* **2009**, *694*, 592. (b) Dupuy, S.; Slawin, A. M. Z.; Nolan, S. P. *Chem.—Eur. J.* **2012**, *18*, 1492.
- (28) (a) Sahoo, B.; Hopkinson, M. N.; Glorious, F. J. *Am. Chem. Soc.* **2013**, *135*, 5505. (b) Shu, X.; Zhang, M.; He, Y.; Frei, H.; Toste, F. D. *J. Am. Chem. Soc.* **2014**, *136*, 5844.